

An aerial photograph of a rural landscape, likely in the Netherlands, showing a patchwork of green and brown fields, a winding river, and a small town. A large white circle is overlaid on the image, centered on the town and river area. The text is overlaid on the top half of the image.

# *Experimentele Technieken*

*in de Hoge Energie Fysica  
en verdere toepassingen*

## Deel 1: Inleiding

Prof. Dr. Albert De Roeck  
CERN, Geneva, Switzerland  
Universiteit Antwerpen  
UC Davis, California



# Kursus inhoud

- Inleiding tot experimenten in de Hoge Energie Fysica
- Voorbereiding op de Large Hadron Collider (LHC).
- Instrumentatie
  - Deeltjes doorheen materie
  - Detector principes
  - Recente ontwikkelingen
- Detail studie van detektoren voor de LHC (CMS, ATLAS, ALICE, LHCb...). Detektor uitdagingen
- Detail studie van andere detektoren (Cosmic rays, lineaire versneller, experimenten voor kosmologie, gravitatiegolven, andere gebieden (geneeskunde...))



# Kursus inhoud

- Inleiding, tot het CMS experiment
- Daarna gespecialiseerde detectoren
- Dan neemt prof D'Hondt over met trigger, DAQ, ....
- Geen examen maar permanente evaluatie
- Opdrachten (1 per persoon). Voorbeelden
  - Deeltjes door materie
  - Detectoren in detail (vanuit leerboeken)
  - CMS Subdetectoren in detail (Tracker, ECAL, HCAL, muons,...)
  - Andere experimenten (WMAP, Auger, Gravitatie experimenten, LHC experimenten)..
  - Echte experimentele papers (van CMS)
    - Start met projecten na de les volgende week



# Projecten

- Voorbeelden van projecten
- Bekijken van detectoren in detail
  - Gas tracker detectoren
  - Halfgeleider tracker detectoren
  - Electromagnetische calorimeters
  - Hadronische calorimeters
  - Deeltjes flow voor precisie metingen
  - Experimentele triggers
  - Speciale experimenten
  - Experimentele metingen en resultaten (CMS/  
Tevatron)



# Timeline & Projects

- Report (5-10 pages) + Presentation of ~ 15 minutes
- When (some time December or January)?
- ⇒ Agenda

**Friday Sept 30- 9:00-12:00 Les 1 (De Roeck) - UA**

**Friday Oct 7 - 9:00-12:00 Les 2**

**Friday Oct 14 – 9:00-12:00 Les 3**

**Friday Nov 18 - 9:00-12:00 Les 4**

**Friday Nov 25 – 9:00-12:00 Les 1 (D'Hondt) - VUB**

**Friday Dec 9 - 9:00-12:00 Les 2**

**Friday Dec 16 - 9:00-12:00 Les 3**

**Friday Dec 23 - 9:00-12:00 Les 4**




# Projects

- Simple Projects (\*)
  - Derive the Bethe Bloch Formula (Gruppen)
  - Cherenkov light radiation (Gruppen)
- Other key experiments (\*\*)
  - The Auger cosmic ray experiment. Ultra High Cosmic Rays
  - The WMAP experiment (satellite, dark matter/dark energy)
  - The Planck experiment (satellite, dark matter dark energy)
  - Ligo: gravitational waves / laser interferometry
  - The MOEDAL experiment at CERN (monopole search)
  - Neutrino experiments (eg OPERA)
- Detection techniques (\*\*)
  - General: EM shower calorimeters (Gruppen)
  - Hadronic shower calorimeters (Gruppen)
  - Compensating calorimeters (Wigmans lectures)
  - Dual readout calorimeters (Wigmans lectures)
  - Silicon detectors (Kleinknecht)
  - MSGC gas detectors
- Difficult projects (\*\*\*)
  - CMS papers with real data (calibration, alignment, efficiency of CMS detector)



# Last Friday: Strange Neutrinos

Faster than the speed of light?



Measure:

- Starting time
- Arrival time
- Distance

Neutrinos arrive 60 nsecs to early





## Why the LHC

The LHC is the most **complex**, most **challenging** and at the same time one of the most **anticipated** scientific instruments so far built by mankind:

The Large Hadron Collider (LHC), built at CERN, Switzerland

- What Is CERN?
- What is the Large Hadron Collider?
- What are the challenges of the collider and experiments?
- What is the science of the Large Hadron Collider?

# CERN

## The European Laboratory for Particle Physics

CERN is the **European Organization for Nuclear Research**, the world's largest Particle Physics Centre, near Geneva, Switzerland  
It is now commonly referred to as **European Laboratory for Particle Physics**  
It was founded in 1954 and has 20 member states + several observer states  
CERN employes **>3000** people + hosts **9000** visitors from **>500** universities.  
Annual budget ~ **1100 MCHF/year** (2010)



CERN: the place where the **World Wide Web** was born



## The 20 CERN Member States



### Member States (Dates of Accession)

 AUSTRIA (1959)	 DENMARK (1953)	 GREECE (1953)	 NORWAY (1953)	 SPAIN (1/1961-12/1968-1/1983)
 BELGIUM (1953)	 FINLAND (1991)	 HUNGARY (1992)	 POLAND (1991)	 SWEDEN (1953)
 BULGARIA (1999)	 FRANCE (1953)	 ITALY (1953)	 PORTUGAL (1986)	 SWITZERLAND (1953)
 CZECH FR (1993)	 GERMANY (1953)	 NETHERLANDS (1953)	 SLOVAK FR (1993)	 UNITED KINGDOM (1953)

CERN AC/DI/MM - E516C 1999 - 15/6/99

US is an observer state... US contributed to the LHC.

**CERN  
Provides  
Particle Beams  
&  
Research Infrastructure**

**Why do we need particle accelerators?**



What is the world made of?  
What holds the world together?  
Where did we come from?






# The BIG BANG

Different types of tools and equipment are needed to observe different sizes of object

Only particle accelerators can explore the tiniest objects in the Universe



The Universe

Many generations of accelerators created  
with higher and higher energy given to the beam particle




Ernest Lawrence  
(1901 - 1958)



CDF

~1500 Scientists

DZero




Tevatron at Fermilab  
 $\times 10^4$  bigger,  $\times 10^6$  higher energy




# Accelerators are Powerful Microscopes.

They make high energy particle beams that allow us to see small things.

$$\lambda = \frac{h}{p}$$



seen by  
low energy beam  
(poorer resolution)



seen by  
high energy beam  
(better resolution)

We can create particles from energy




Two beams of protons collide and generate, in a very tiny space, temperatures over a billion times higher than those prevailing at the center of the Sun.

# The Fundamental Forces of Nature


**Electromagnetism:**  
gives light, radio, holds atoms together

**Strong Nuclear Force:**  
holds nuclei together

**Weak Nuclear Force:**  
gives radioactivity



together  
they make  
the Sun  
shine



**Gravity:**  
holds planets and stars together





# The Standard Model in Particle Physics


But not all questions solved:

Why is the top quark much more heavy than the quarks  
 $\Rightarrow$  Mass(top) = gold nucleus

What is the origin of mass?


Astrophysics/cosmological measurements show that most matter in the universe is **NOT** in this table

What is this Dark Matter?




Four known forces

- Gravity
- Electro-magnetism
- Strong nuclear force
- Weak force



quarks


proton



gluons

neutron

Create (anti)particles that existed  
~0.001 nanosecond after Big Bang



Accelerators

Create (anti)particles that existed  
~0.001 nanosecond after Big Bang

$$E = mc^2$$

LHC

Inflation

Big Bang

One Force

Four Forces

particles  
anti-particles


possible da  
tter relicts

cosmic micro  
radiation visible

Today

particles

*Do all the forces become one?  
Extra hidden dimensions of space?  
Where did all antimatter go?*






1. Are there undiscovered principles of nature:  
New symmetries, new physical laws?
2. How can we solve the mystery of dark energy?
3. Are there extra dimensions of space?
4. Do all the forces become one?
5. Why are there so many kinds of particles?
6. What is dark matter?  
How can we make it in the laboratory?
7. What are neutrinos telling us?
8. How did the universe come to be?
9. What happened to the antimatter?
10. What is mass?

“Quantum Universe” and  
“Discovering the Quantum Universe”

Discoveries and breakthroughs will likely come from  
Energy Frontier Accelerators at the Terascale.


**Evolved Thinker**





# The LHC = a proton proton collider

7 TeV + 7 TeV




1 TeV = 1 Tera electron volt  
=  $10^{12}$  electron volt

## Primary physics targets

- Origin of mass
- Nature of Dark Matter
- Understanding space time
- Matter versus antimatter
- Primordial plasma

The LHC will determine the Future course of High Energy Physics  
The LHC has started with collisions in 2009

# Accelerators for Charged Particles



- A** Cathode
- B** Conductive coating
- C** Anode
- D** Phosphor-coated screen
- E** Electron beams
- F** Shadow mask





Photo courtesy SLAC  
Schematic diagram of a cyclotron





# Recent High Energy Colliders

Highest energies can be reached with proton colliders

Machine	Year	Beams	Energy ( $\sqrt{s}$ )	Luminosity
SPPS (CERN)	1981	pp	630-900 GeV	$6 \cdot 10^{30} \text{cm}^{-2} \text{s}^{-1}$
Tevatron (FNAL)	1987	pp	1800-2000 GeV	$10^{31}-10^{32} \text{cm}^{-2} \text{s}^{-1}$
SLC (SLAC)	1989	$e^+e^-$	90 GeV	$10^{30} \text{cm}^{-2} \text{s}^{-1}$
LEP (CERN)	1989	$e^+e^-$	90-200 GeV	$10^{31}-10^{32} \text{cm}^{-2} \text{s}^{-1}$
HERA (DESY)	1992	ep	300 GeV	$10^{31}-10^{32} \text{cm}^{-2} \text{s}^{-1}$
RHIC (BNL)	2000	pp / AA	200-500 GeV	$10^{32} \text{cm}^{-2} \text{s}^{-1}$
<b>LHC (CERN)</b>	<b>2009</b>	<b>pp (AA)</b>	<b>10-14 TeV</b>	<b><math>10^{33}-10^{34} \text{cm}^{-2} \text{s}^{-1}</math></b>

Luminosity = number of events/cross section/sec

- Limits on circular machines
  - Proton colliders: Dipole magnet strength  $\rightarrow$  superconducting magnets
  - Electron colliders: Synchrotron radiation/RF power

**Accelerators:** developed in physics labs  
are used in hospitals for HADRON Therapy



Around 9000 of the 17000 accelerators operating in the world  
today are used for medicine

# The LHC: >20 Years Already!


## LHC History

- 1982 : First studies for the LHC project
- 1983 : Z0/W discovered at SPS proton antiproton collider
- 1989 : Start of LEP operation (Z boson-factory)
- 1994 : Approval of the LHC by the CERN Council
- 1996 : Final decision to start the LHC construction
- 1996 : LEP operation > 80 GeV (W boson -factory)
- 2000 : Last year of LEP operation above 100 GeV
- 2002 : LEP equipment removed
- 2003 : Start of the LHC installation
- 2005 : Start of LHC hardware commissioning
- 2008 : Expected LHC commissioning with beam

Luminosity=# events/cross section/sec

1984

ECFA 84/85  
CERN 84-10  
5 September 1984



1984: cms energy	10-18 TeV
Luminosity	$10^{31}-10^{33}\text{cm}^{-2}\text{s}^{-1}$
1987: cms energy	16 TeV
Luminosity	$10^{33}-10^{34}\text{cm}^{-2}\text{s}^{-1}$
Final: cms energy	14 TeV
Luminosity	$10^{33}-10^{34}\text{cm}^{-2}\text{s}^{-1}$




 SEARCH


## TIME's Best Inventions of 2008

### The Other 49 Best Inventions


◀ Previous    Next ▶

### 5. The Large Hadron Collider

If someone invented a practical 200-m.p.g. automobile and that automobile got a flat tire, nobody would claim that the car itself was a failure. The same applies to the Large Hadron Collider, the world's biggest particle accelerator, which went online in September, ran for 10 days and then had to shut down at least until next spring because of an overheated wire. The mammoth machine will send protons wheeling in opposite directions at nearly the speed of light, then smash them together at 6,000 times a second to try to answer such deep questions as why mass exists and whether the universe has extra dimensions. If it takes a few extra months to find out, so what?

ARTICLE TOOLS

- Print
- Email
- Sphere
- AddThis
- RSS
- Yahoo! Buzz



STEFANO DAL POZZOLO / CONTRASTO / REDUX

#### Top Stories on Time.com

- The Fight Over Gay Adoption Heats Up
- Bush's Last Days: The Lamest Duck
- Thailand's Political Crisis Becomes a Global One
- The Meaning of Paul Volcker's Comeback
- Papua Proposal: A Microchip to Track the HIV-Positive

# The LHC Machine and Experiments

LHC is 100m underground




LHC is 27 km long

Magnet Temperature is 1.9 Kelvin = -271 Celsius

LHC has ~ 9000 magnets

LHC: 40 million proton-proton collisions per second


LHC: Luminosity  $10\text{-}100 \text{ fb}^{-1}/\text{year}$  (after start-up phase)



- High Energy  $\Rightarrow$  factor 7 increase w.r.t. present accelerators
- High Luminosity (# events/cross section/time)  $\Rightarrow$  factor 100 increase

# The Cryodipole Magnets

- Superconducting (1.9 K) dipoles producing a field of 8.4 T
- Cost: ~ 0.5 million CHF each. We need 1232 of them
- **Total stored energy in all magnets = 11GJ**  
= energy of an Airbus 380 flying at 700 km/hour
- One dipole weighs around 34000 kg





# LHC RF system

- ❑ The LHC RF system operates at 400 MHz.
- ❑ It is composed of 16 superconducting cavities, 8 per beam.
- ❑ Peak accelerating voltage of 16 MV/beam.

For LEP at 104 GeV : 3600 MV/beam !

	Synchrotron radiation loss
LHC @ 7 TeV	6.7 keV /turn
LEP @ 104 GeV	~3 GeV /turn


The LHC beam radiates a sufficient amount of visible photons to be actually observable with a camera !  
(total power ~ 0.2 W/m)



LHC facts

100 m underground: a tunnel 27 km circumference

One of the **fastest** racetracks on the planet



$v \approx 27\text{km/h}$   
1turn in 1h





Several thousand billion protons traveling at 99.9999991% of the speed of light will travel round the 27km ring over 11000 times a second



# LHC facts

The **emptiest** space in the solar system...



To accelerate protons to almost the speed of light, we need a vacuum similar to interplanetary space. The pressure in the beam-pipes of the LHC will be about ten times lower than on the moon.



# LHC facts


One of the **coldest** places in the Universe...

the largest cryogenic system ever built  
54 km fridge!




With a temperature of around -271 degrees Celsius, or 1.9 degrees above absolute zero, the LHC is colder than interstellar space.

One of the **hottest** places in the Galaxy...



Simulation of a collision in the CMS experiment



Simulation of a collision in the ALICE experiment

When two beams of protons collide, they generate within a tiny volume, temperatures more than a billion times those in the very heart of the Sun.



# Energy in the beam


## Comparison...

The energy of an A380 at 700 km/hour corresponds to the energy stored in the LHC magnet system :  
Sufficient to heat up and melt 12 tons of Copper!!



The energy stored in one LHC beam corresponds approximately to...



- 90 kg of TNT
- 8 litres of gasoline
- 15 kg of chocolate




It's how ease the energy is released that matters most !!



# The Accelerator Scheme




# CERN's particle accelerator chain




Run Number: 177531,  
Event Number: 183764  
Date: 2011-03-13, 18:20:50 CET

Cells: Tiles, EMC, EMEC, HEC

LHC Large Hadron Collider    SPS Super Proton Synchrotron    PS Proton Synchrotron  
AD Antiproton Decelerator    CTF3 Clic Test Facility    CNRS Cern Neutrinos to Gran Sasso    ISOLDE Isotope Separator OnLine Device  
LER Low Energy Ion Ring    LINAC LINear ACcelerator    n-ToF Neutrons Time Of Flight







# Experiments at the LHC








# The Four Main LHC Experiments






# Detectors are essentially completed



Atlas & CMS construction started 9 years ago





+TOTEM, LHCf, MOEDAL

Now gearing up for first collisions...





# LHC Story: Beam Halo and Splashes on 10/9/08







First LHC activity in the detectors in Sept. 2008, followed by the LHC accident

# LHC Story: First Collisions in Nov. 2009

23/11 First 'trial' collisions in the experiments

A run with collisions at  $\sqrt{s} = 900$  GeV or 2.36 TeV in December 2009






# Key Moments




Some of the  
key moments  
the last years




# 30/3/10: Experiments are waiting for 7 TeV...



12:58  
7 TeV collisions!!!




# First Collisions at 7 TeV




A Brave New World....





Length = 55 m    Width = 32 m    Height = 35 m    but spatial precision ~ 100  $\mu\text{m}$

# The Compact Muon Solenoid Experiment



In total about

~100 000 000 electronic channels

Each channel checked

40 000 000 times per second (collision rate is 40 MHz)

An on-line trigger selects events and reduces the rate from 40MHz to 100 Hz

Amount of data of just one collisions


>1 500 000 Bytes



# Detectors at Accelerators

Particle Detection: What we “see” as particles:


For “stable particles” of life time  $\geq 10^{-10}$  s:



For charged tracks :  $\Delta p/p \propto p,$

for calorimetry :  $\Delta E/E \propto \frac{1}{\sqrt{E}}.$

# Particles in the detector




Hoe werken deze detektoren?  
⇒ Lessen 2,3,4




# The LHC Detectors are Major Challenges

- CMS/ATLAS detectors have about 100 million read-out channels
- Collisions in the detectors happen every 25 nanoseconds
- ATLAS uses over 3000 km of cables in the experiment
- The data volume recorded at the front-end in CMS is 1 TB/second which is equivalent to the world wide communication network traffic
- Data recorded during the 10-20 years of LHC life will be about all the words spoken by mankind since its appearance on earth
- A worry for the detectors: the kinetic energy of the beam is that of a small aircraft carrier of  $10^4$  tons going 20 miles/ hour



ATLAS pixel detector

>200 m<sup>2</sup> of silicon



CMS silicon tracker



Object	Weight (tons)
Boeing 747 [fully loaded]	200
Endeavor space shuttle	368
ATLAS	7,000
Eiffel Tower	7,300
USS John McCain	8,300
CMS	12,500

# CMS Solenoid

The largest high field solenoid magnet ever build!!

Successfully tested in August '06!!

Magnetic length	12.5 m
Free bore diameter	6 m
Central magnetic induction	4 T $\approx 100,000$ times earth magnetic field
Temperature	4.2 degrees Kelvin $\approx -269$ degrees Celcius
Nominal current	20 kA
Stored energy	2.7 GJ
Magnetic Radial Pressure	64 Atmospheres



CMS


ALEPH

factor



# Construction of CMS ( $\geq 2002$ )

...In a large hall on surface



**Muon detectors**  
832 000 read-out channels





# Lowering of the Solenoid

The Central piece of CMS  
⇒ The barrel wheel with the solenoid

Total weight ~ 2Ktons  
= 5 jumbo jets

Lowered February 28 (2007)






# Lowering of CMS in the Underground Cavern

The first force CMS carefully studied was.... Gravity






CMS built on the surface and lowered in the cavern 100m below  
Piece by piece over three years



Hydraulic jacks and control tower used in CMS will be used  
in Durban to lift the roof of the stadium for World Cup 2010







# The CMS Detector: Calorimeters

ECAL: Barrel 36 super modules/1700 crystals  
Endcaps detectors completed in summer 2008  
Total of ~70000 crystals for this detector



Lead tungstate.  
Transparent like glass  
Heavy as lead!!




Central ECAL installation in CMS

Hadronic Calorimeter (brass/scintillator)  
completed in 2006  
Lowering in the experimental hall





# Hadron Calorimeter (HCAL)



Made of dense brass layers interspersed with plastic scintillators

Used over a million World War II brass shell casements from the Russian Navy in making some of its detector components; is made up of 36 wedges, each of which weighs as much as 6 African elephants; contains over 400 “optical decoder” units, all of which were made by American high school students through the QuarkNet programme.

## ***Function:***



Measure energy of hadrons (protons, neutrons)







# The CMS Central Tracker



- 200 m<sup>2</sup> silicon detectors (~ tennis court)
- ~ 10<sup>7</sup> read-out channels: silicon strips




# Installation of the Central Tracker in CMS

December 2007






# Pixel Tracking Detector



In total  $7 \cdot 10^7$  read out channels  
~ photo camera with 70 million pixels  
taking 40 million photos per second!!







# CMS Works! ...

## Example: Recorded Cosmic Muons

Muons recorded by a part of CMS  
(2-5%) during the summer of 2006



Muons recorded by the complete  
central tracker during the summer '08



# The LHC Data Challenge

The LHC accelerator will run for 10-15 years

Experiments will produce about **15 Million Gigabytes** of data each year (about 20 million CDs!)


LHC data analysis requires a computing power equivalent to **~100,000 of today's fastest PC processors**

Requires many cooperating computer centres, as CERN can only provide **~20% of the capacity**





The LHC computing challenge  
**furthest reaching** computer in the world...



Tens of thousands of computers based all over the world are used to analyse data from CERN. The computing GRID is the next advance in decentralised computing from the laboratory that brought you the World-Wide Web.

# LHC Computing Grid project (LCG)


More than 140 computing centres  
12 large centres for primary data  
management:  
CERN (Tier-0)

Eleven Tier-1s

38 federations of smaller Tier-2  
centres

India – BARC, TIFR, VECC

35 countries involved





# The Science of the LHC

⇒ Explore the new high energy regime: The Terascale

# The Origin of Mass

Some particles have mass, some do not

Where do the masses  
come from ?

Newton:



Weight proportional to Mass

Einstein:

Energy related to Mass

Neither explained origin of Mass

Explanation of Profs P. Higgs  
R. Brout en F. Englert  
⇒ A new field and particle








# The Higgs Particle

⇒ What is the **origin of mass** of the elementary particles?  
Solution within the Standard Model: A scalar Higgs field



⇒ At least one new scalar particle should exist: The Higgs  
The Higgs is the **last missing particle in the Standard Model**  
**One of the main missions of LHC: discover the Higgs**



- If the Higgs exist: LHC will discover it after 2-3 years of operation
- If the Higgs does not exist: LHC should see other spectacular new effects

# Beyond the Higgs Particle


## Supersymmetry: a new symmetry in Nature



Candidate particles for Dark Matter  
 $\Rightarrow$  Produce Dark Matter in the lab

SUSY particle production at the LHC


+ 2 D-jets  
 + 4 jets





# Detecting Supersymmetric Particles

Energy produced in the detector





Supersymmetric particles decay and produce a cascade of jets, leptons and missing (transverse) energy due to escaping 'dark matter' particles

➔ Very clear signatures in CMS and ATLAS

LHC can discover supersymmetric partners of the quarks and gluons as heavy as 2 to 3 TeV

The expected cross sections are huge!!  $\Rightarrow$  10,000 to 100,000 particles per year

# Supersymmetry: a new symmetry in Nature




Susy partners are very heavy  
 $\sim 1000$  mass of the proton



# Sparticle Reconstruction

Mass precision for a favorable benchmark point at the LHC  
LCC1~ SPS1a~ point B'


$m_0=100 \text{ GeV}$   
 $m_{1/2}= 250 \text{ GeV}$   
 $A_0=-100$   
 $\tan\beta = 10$   
 $\text{sign}(\mu)=+$




hep-ph/0508198

Lightest neutralino  $\rightarrow$  Dark Matter?  
Fit SUSY model parameters to the measured SUSY particle masses to extract  $\Omega_\chi h^2 \Rightarrow O(10\%)$

GeV	LHC
$\Delta m_{\tilde{\chi}_1^0}$	4.8
$\Delta m_{\tilde{\chi}_2^0}$	4.7
$\Delta m_{\tilde{\chi}_4^0}$	5.1
$\Delta m_{\tilde{t}_R}$	4.8
$\Delta m_{\tilde{\ell}_L}$	5.0
$\Delta m_{\tau_1}$	5-8
$\Delta m_{\tilde{q}_L}$	8.7
$\Delta m_{\tilde{q}_R}$	7-12
$\Delta m_{\tilde{b}_1}$	7.5
$\Delta m_{\tilde{b}_2}$	7.9
$\Delta m_{\tilde{g}}$	8.0



D. Miller et al  
 $\Rightarrow$  Use shapes






# Beyond the Higgs Particle

## Extra Space Dimensions

Problem:

$$m_{EW} = \frac{1}{(G_F \cdot \sqrt{2})^{1/2}} = 246 \text{ GeV}$$

$$M_{Pl} = \frac{1}{\sqrt{G_N}} = 1.2 \cdot 10^{19} \text{ GeV}$$





The Gravity force becomes strong!



# Detecting Extra Dimensions at the LHC

Main detection modes at the experiments

- Large missing (transverse) energy
- Resonance production





LHC can detect extra dimensions for scales up to 5 to 9 TeV

# Quantum Black Holes at the LHC?

Black Holes are a direct prediction of Einstein's general theory on relativity

If the Planck scale is in  $\sim$ TeV region:  
can expect Quantum Black Hole production



Simulation of a Quantum Black Hole event

Quantum Black Holes are harmless for the environment: they will decay within less than  $10^{-27}$  seconds

Quantum Black Holes open the exciting perspective to study Quantum Gravity in the lab!



# Quantum Back Holes

- Schwarzschild radius

4-dim.,  $M_{\text{gravity}} = M_{\text{Planck}}$

4 + n-dim.,  $M_{\text{gravity}} = M_D \sim \text{TeV}$



Since  $M_D$  is low, tiny black holes of  $M_{\text{BH}} \sim \text{TeV}$  can be produced if partons  $ij$  with  $\sqrt{s_{ij}} = M_{\text{BH}}$  pass at a distance smaller than  $R_s$

- Large partonic cross-section:  $\sigma(ij \rightarrow \text{BH}) \sim \pi R_s$
- $\sigma(pp \rightarrow \text{BH})$  is in the range of 1 nb - 1 fb

e.g. For  $M_D \sim 1 \text{ TeV}$  and  $n=3$ , produce 1 event/second at the LHC

- Black holes decay immediately by Hawking radiation (democratic evaporation):


- large multiplicity
- small missing  $E$
- jets/leptons  $\sim 5$

expected signature (quite spectacular ...)

Landsberg, Dimopoulos  
Giddings, Thomas, Rizzo...

$$R_s \rightarrow \ll 10^{-35} \text{ m}$$

$$R_s \rightarrow \sim 10^{-19} \text{ m}$$



# Quantum Black Holes

- Can LHC destroy the planet?  
⇒ **No!**
- See the report of the LHC Safety assesment group (LSAG) <http://arXiv.org/pdf/0806.3414>
- More information on
  - S.B. Giddings and M. Mangano, <http://arXiv.org/pdf/0806.3381> LSAG, <http://arXiv.org/pdf/0806.3414>
  - Scientific Policy Committee Review, <http://indico.cern.ch/getFile.py/access?contribId=20&resId=0&materialId=0&confId=35065>
  - CERN public web page, <http://public.web.cern.ch/public/en/LHC/Safety-en.html>






# Black Holes Hunters at the LHC...



## Scientific American


### THE RISE AND DEMISE OF A QUANTUM BLACK HOLE






# New Physics at the LHC


## New Gauge Bosons?




## Hidden Valleys?




## ZZ/WW resonances?




## Technicolor?




## Extra Dimensions?




## Black Holes???



## Little Higgs?




## Split Susy?



We do not know what is out there for us...



# LHC Luminosity/Sensitivity with Time






F. Moortgat, A. De Roeck

First physics run:  $O(1\text{fb}^{-1})$

Before LHC incident: add one year now

# Matter-Antimatter



The properties and subtle differences of matter and anti-matter using mesons containing the beauty quark, will be studied further in the **LHCb experiment**





# Primordial Plasma

Lead-lead collisions at the LHC to study the primordial plasma, a state of matter in the early moments of the Universe




Study the phase transition of a state of **quark gluon plasma** created at the time of the early Universe to the **baryonic matter** we observe today

A lead lead collision simulated in the ALICE detector


# A Recorded Heavy Ion Collision

CMS Experiment at the LHC, CERN  
Mon 2010-Nov-08 11:22:07 CET  
Run 150431 Event 541464  
C.O.M. Energy 7Z TeV





# The big Bang



Electro-weak phase transition (ATLAS, CMS...)

QCD phase transition (ALICE...)




LHC will study the first 10<sup>-10</sup> - 10<sup>-5</sup> seconds...



# Other Detectors



Dark matter/dark energy



WMAP 5-year  
-200  $T(\mu\text{K})$  +200




# CERN and Technology

⇒ Direct Spin-off of the technologies developed and used at CERN

# Applications of Grid Computing

Multitude of applications from a growing number of domains


- Archeology
- Astronomy & Astrophysics
- Civil Protection
- Computational Chemistry
- Earth Sciences
- Financial Simulation
- Fusion
- Geophysics
- High Energy Physics
- Life Sciences
- Multimedia
- Material Sciences
- ...




**Infrastructure used by >5000 researchers  
- submitted ~20 millions jobs in 2006**




# Technology Transfer Projects




Silicon detector for a Compton camera in nuclear medical imaging




Radio-isotope production for medical applications



Thin films by sputtering or evaporation





Medipix: Medical X-ray diagnosis with contrast enhancement and dose reduction




Radiography of a bat, recorded with a GEM detector


# Fundamental research has always been a driving force for innovation



Relativity  
100%  
SCIENCE



For GPS to work, we have to take into account the correction due to time dilation. Otherwise, there would be a position error of around 10m after just 5 minutes of travel-time!



Electromagnetism

100%  
SCIENCE



Telephones use electromagnetic waves to communicate




# Have you heard of the World Wide Web or the Internet ?




**Tim Berners Lee  
Developed the  
WWW at CERN  
initially for  
sharing particle  
physicists data**

**I think there is a world market  
for maybe five computers.  
--THOMAS WATSON, chairman of  
IBM, 1943.**



# Instrumentation at CERN





Detectors used in LHC


We use state-of-the-art instruments to explore our understanding of nature. Physicists take this knowledge to other fields. Studies have demonstrated that the transfer of knowledge from fundamental research enables high-tech companies to remain on the cutting edge of innovation and generates a variety of social and economic benefits. It also has an important impact on our culture and education.



# Detectors: developed in physics labs are used for medical imagery



PET (Positron Emission Tomography) is a very important technique for localising and studying certain types of cancer using the Fluor-18 isotope produced by particle accelerators. PET uses antimatter (positrons).



In the recent report from the Intergovernmental Panel on Climate Change, data from various models and sources were combined to project the future climate. This image shows Scenario A1B: simulated mean temperature change relative to 1980-1999.

## Scientific Applications:

### EGEE Makes Rapid Earthquake Analysis Possible

Using the advanced Grid infrastructure of the Enabling Grids for E-science (EGEE) project, researchers at the Institut de Physique du Globe de Paris (IPGP), France, were able to analyze, within 30 hours of it occurring, the large Indonesian earthquake that struck on March 28. Although less severe than the one in December, which caused a tsunami wave in the Indian Ocean, more than 1,000 people were killed in this second major earthquake.

The analysis showed that the March earthquake was not a belated aftershock of the December one, although they are intricately linked. The March earthquake was probably triggered by the one in December, but happened in a different part of the fault line further south, and the mechanisms of the two earthquakes were different. Although the basic geometry of the region is known, the strength of the earthquake was astonishing.

Understanding the exact parameters of when, where and how an earthquake occurs brings researchers closer to comprehending why earthquakes happen. This might make it possible to predict when and where earthquakes will happen in the future and to assess the potential impact they could have on specific regions. Rapid analysis is particularly important for the relief efforts after a major earthquake, where those in charge need to have accurate information about the epicenter, magnitude

## Grids open new perspectives to *in silico* drug discovery

- Reduced cost and adding an accelerating factor in the search for new drugs
- Diseases such as HIV/AIDS, SRAS, Bird Flu etc. are a threat to public health due to world wide exchanges and circulation of people



- Home
- News
- Markets
- Personal Finance
- Infotech
- Opinion
- News By Company
- News By Industry
- Economy
- International Business
- Auto
- Banking/ Finance
- Cons. Products
- Energy
- Ind'l Goods / Svs
- Healthcare

## Scientists gear up to tackle 15 million gigabytes of data

4 Nov 2008, 16:25 hrs IST, IANS

Print | EMail | Discuss | Share | Save | Comment | Text: ( )

LONDON: The four huge detectors of the new Large Hadron Collider near Geneva, when fully operational, are expected to generate up to a staggering 15 million gigabytes of data every year.

**Vostro™ 1310 Laptop**

Intel™ Core™ 2 Duo Processor T5670

Genuine Windows Vista® Home Basic

1GB SDRAM + 160GB HDD

13.3" Widescreen Display

Now Only Rs. 40,900\*\*

Costs more Now

Andreas Hirstius, manager of Openlab and the CERN Scientific Computing, explained how computer scientists have met the challenge of handling this unprecedented volume of data.

When CERN staff first considered the mid-1990s how they might cope with the large volume of data produced when its two beam proton colliders collide, a single gigabyte of disk space still cost a few hundred dollars and CERN's total connectivity was equivalent to just

of today's [broadband connections](#).

It quickly became clear that computing power at CERN, even taking Moore's Law into account, would be significantly less than that required to analyse Large Hadron Collider (LHC) data.

The solution, which transpired during the 1990s, was to turn to "high-throughput computing" where the focus is not on shifting data as quickly as possible from one place to another but rather from shifting as much information as possible between those two places.

[High-performance computing](#) is ideal for particle physics because the data produced in the millions of proton-proton collisions are all independent of one another - and therefore can be handled independently, according to an Institute of Physics (IOP) report.

- NEWS
- COMMENT
- BUSINESS
- MONEY
- SPORT
- LIFE & STYLE
- TRAVEL
- DRIVING
- ARTS
- UK NEWS
- WORLD NEWS
- POLITICS
- ENVIRONMENT
- WEATHER
- TECH & WEB
- VIDEO
- PHOTO

Where am I? > Home > News > UK News > Science News

From [The Sunday Times](#)

September 7, 2008

## Large Hadron Collider could help fight cancer

### Stand by for advances in health and climate research as the 'big bang' machine starts up

Jonathan Leake

The giant new particle collider at Europe's centre for nuclear research, which is due to start work on Wednesday, is being linked to spectacular spin-offs including improved cancer treatments, systems for destroying nuclear waste and insights into climate change.

"Everyone is looking at the start up of the Large Hadron Collider [LHC] but Cern has many other research programmes with important practical uses," said Paul Collier, who runs the main control room at the European Organisation for Nuclear Research (Cern).

The first beams of particles have been successfully fired around nearly half of the 17-mile tunnel in Switzerland, where Cern is based. Linked research has already spurred useful byproducts.

In a typical year, the huge machine, which will smash particles into each other at enormous speed, should generate enough data to fill 56m CDs. That means physicists have had to create a sophisticated system for organising information extremely quickly. The Grid, as they call it, is likely to become the model for many other systems designed to handle large volumes of data.

Another project has suggested a potentially radical new way of dealing with nuclear waste. Cern's physicists found that firing a beam of protons (a type of sub-atomic particle) into blocks of lead could

#### RELATED LINKS

- > LHC will not turn world to goo - scientists
- > How the press demeans

#### TIMES RECOMMENDS

- > Spike Milligan has the last laugh
- > Pubs told to bring an end to happy hours
- > Pianist's bequeathed skull stars in Hamlet

#### PARENT POWER



#### Britain's best schools


The Sunday Times Parent Power: the UK's top schools ranked by the latest examination results

Archeology  
Astronomy  
Astrophysics  
Civil Protection  
Comp. Chemistry  
Earth Sciences  
Finance  
Fusion  
Geophysics  
High Energy Physics  
Life Sciences  
Multimedia  
Material Sciences  
...

Scheduled = 21539  
Running = 25374

>250 sites  
48 countries  
>50,000 CPUs  
>20 PetaBytes  
>10,000 users  
>150 VOs  
>150,000 jobs/day

**21:13:50 UTC**







# CERN as an Educator



**The End**




# Mogelijke Projecten

Performance of the CMS Pixel detector with cosmic ray data	 CFT-09-001	
<i>Performance of the CMS Pixel detector with cosmic ray data</i>		
The CMS Silicon Strip Tracker Operation and Performance With Cosm ...	 CFT-09-002	
<i>The CMS Silicon Strip Tracker Operation and Performance With Cosmic Rays in 3.8 T Magnetic</i>		
Alignment of the CMS Silicon Tracker During Commissioning with Co ...	 CFT-09-003	
<i>Alignment of the CMS Tracker with CRAFT data</i>		
Performance and Operation of the CMS Crystal Electromagnetic Calo ...	 CFT-09-004	
<i>Performance and Operation of the CMS Crystal Electromagnetic Calorimeter</i>		
Measurement of the muon stopping power in PbWO	 CFT-09-005	
Time reconstruction and performance of Crystal ECAL	 CFT-09-006	
From Detector to Analysis: CMS Data Processing Workflows During a ...	 CFT-09-007	
The CMS CRAFT Exercise	 CFT-09-008	
<i>Overview of the CRAFT exercise</i>		
Performance of CMS Hadron Calorimeter Using Cosmic Muons and the ...	 CFT-09-009	
<i>Performance of CMS Hadron Calorimeter Using Cosmic Muons and the Proton Beam From the</i>		
Measurements of RPC performance in CMS with cosmic rays	 CFT-09-010	
CSC Performance	 CFT-09-011	
<i>Summary of the performance of the CSC's from CRAFT Data</i>		
Results On Local Muon Reconstruction in DT Chambers From Analysis ...	 CFT-09-012	
<i>Results On Local Muon Reconstruction in DT Chambers From Analysis of Cosmic Muon Data</i>		
L1 Trigger Performance	 CFT-09-013	
<i>Performance of triggers run in CRAFT</i>		
Muon Reconstruction Performance	 CFT-09-014	
Magnetic Field Studies	 CFT-09-015	
		Muon Track-based alignment
		 CFT-09-016
		<i>Alignment of the CMS Muon System with Cosmic Ray and Beam-Halo Tracks   Samples: 2_2</i>
		Muon hardware-based alignment
		 CFT-09-017
		HCAL Timing
		 CFT-09-018
		<i>Presentation of the HCAL Timing and Synchronization system, validation results from CRAFT a</i>
		Anomalous signals in HCAL
		 CFT-09-019
		<i>Describe performance of algorithms designed to flag bad or problematic cells in HCAL   Sample</i>
		Performance of the High Level Trigger
		 CFT-09-020
		Characterization of Beam Halo Data
		 CFT-09-021
		DT Local Trigger Performance
		 CFT-09-022
		<i>Performances of the CMS Drift-Tubes Local Trigger during CRAFT</i>
		DT Calibration
		 CFT-09-023
		<i>Results on the DT Calibration and Drift Velocity analysis with CRAFT data</i>
		Craft Test Paper
		 CFT-09-024
		<i>This is a dummy paper to test the various steps in publication of CMS papers (Remarks: ignore this</i>
		Fine Synchronization of the muon Drift-Tubes local trigger
		 CFT-09-025
		<i>Fine synchronization of the DT local trigger on CRAFT reported and method for LHC presented</i>

# Integrated Luminosity @ 7 TeV

CMS values    Similar ones for ATLAS



Since end of March (7 TeV):  
14.5 pb<sup>-1</sup> delivered (\*)  
13.2 pb<sup>-1</sup> recorded (\*)  
Data taking efficiency (> 90%)


(\*) Stable beams only

- Max Lumi now  $\sim 6.8 \cdot 10^{31} \text{ cm}^{-2} \text{ s}^{-1}$
- Aim for this year  $10^{32} \text{ cm}^{-2} \text{ s}^{-1}$ ...

Luminosity = number of events  
per cross section per second



# Last week's running...



$1 \cdot 10^{31} \text{ cm}^{-2}\text{s}^{-1}$  instantaneous luminosity reached!

Only a factor 10 to go to the end-year goal.

⇒ Bunch trains! Getting > 300 colliding bunches in CMS/ATLAS

# Last days' running...

## PROTON PHYSICS: STABLE BEAMS

Energy:

3500 GeV


I(B1):

2.12e+13


I(B2):

2.00e+13

FBCT Intensity and Beam Energy Updated: 09:03:58



Instantaneous Luminosity Updated: 09:03:58



$6.8 \cdot 10^{31} \text{ cm}^{-2}\text{s}^{-1}$  instantaneous luminosity reached! (200 bunches)

Only a factor 1.5 to go to the end-year goal.

⇒ Bunch trains! Getting > 300 colliding bunches in CMS/ATLAS